

SPECIALTY GUIDELINE MANAGEMENT

SIMPONI (golimumab for subcutaneous injection)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

A. FDA-Approved Indications

1. Moderately to severely active rheumatoid arthritis (RA) in combination with methotrexate
2. Active psoriatic arthritis (PsA)
3. Active ankylosing spondylitis (AS)
4. Moderately to severely active ulcerative colitis (UC)

B. Compendial Use

Axial spondyloarthritis

All other indications are considered experimental/investigational and not medically necessary.

II. DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review:

A. Rheumatoid arthritis (RA)

1. For initial requests:
 - i. Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
 - ii. Laboratory results, chart notes, or medical record documentation of biomarker testing (i.e., rheumatoid factor [RF], anti-cyclic citrullinated peptide [anti-CCP], and C-reactive protein [CRP] and/or erythrocyte sedimentation rate [ESR]) (if applicable).
2. For continuation requests: Chart notes or medical record documentation supporting positive clinical response.

B. Psoriatic arthritis: For continuation requests: Chart notes or medical record documentation supporting positive clinical response.

C. Ankylosing spondylitis (AS) and axial spondyloarthritis

1. Initial requests: Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable). If therapy is not advisable, documentation of clinical reason to avoid therapy.
2. Continuation requests: Chart notes or medical record documentation supporting positive clinical response.

D. Ulcerative colitis (UC)

1. Initial requests: Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
2. Continuation requests: Chart notes or medical record documentation supporting positive clinical response to therapy or remission.

III. CRITERIA FOR INITIAL APPROVAL

A. Moderately to severely active rheumatoid arthritis (RA)

1. Authorization of 12 months may be granted for members who have previously received a biologic or targeted synthetic DMARD (e.g., Rinvoq, Xeljanz) indicated for moderately to severely active rheumatoid arthritis. The requested medication must be prescribed in combination with methotrexate unless the member has a clinical reason not to use methotrexate (see Appendix A).
2. Authorization of 12 months may be granted for treatment of moderately to severely active RA when all of the following criteria are met:
 - i. Member meets either of the following criteria:
 - a. Member has been tested for either of the following biomarkers and the test was positive:
 1. Rheumatoid factor (RF)
 2. Anti-cyclic citrullinated peptide (anti-CCP)
 - b. Member has been tested for ALL of the following biomarkers:
 1. RF
 2. Anti-CCP
 3. C-reactive protein (CRP) and/or erythrocyte sedimentation rate (ESR)
 - ii. Member is prescribed the requested medication in combination with methotrexate or has a clinical reason not to use methotrexate.
 - iii. Member meets either of the following criteria:
 - a. Member has experienced an inadequate response to at least a 3-month trial of methotrexate despite adequate dosing (i.e., titrated to at least 15 mg/week).
 - b. Member has an intolerance or contraindication to methotrexate (see Appendix A).

B. Active psoriatic arthritis (PsA)

Authorization of 12 months may be granted for treatment of active psoriatic arthritis (PsA).

C. Active ankylosing spondylitis (AS) and active axial spondyloarthritis

1. Authorization of 12 months may be granted for members who have previously received a biologic indicated for active ankylosing spondylitis or active axial spondyloarthritis.
2. Authorization of 12 months may be granted for treatment of active ankylosing spondylitis and active axial spondyloarthritis when any of the following criteria is met:
 - i. Member has experienced an inadequate response to at least two non-steroidal anti-inflammatory drugs (NSAIDs).
 - ii. Member has an intolerance or contraindication to two or more NSAIDs.

D. Moderately to severely active ulcerative colitis (UC)

1. Authorization of 12 months may be granted for members who have previously received a biologic or targeted synthetic drug (e.g., Xeljanz) indicated for moderately to severely active ulcerative colitis.
2. Authorization of 12 months may be granted for the treatment of moderately to severely active UC when the member has had an inadequate response, intolerance or contraindication to at least one conventional therapy option (see Appendix B).

IV. CONTINUATION OF THERAPY

A. Moderately to severely active rheumatoid arthritis (RA)

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active rheumatoid arthritis and who achieve or maintain a positive clinical response as evidenced by disease activity improvement of at least 20% from baseline in tender joint count, swollen joint count, pain, or disability.

B. Active psoriatic arthritis (PsA)

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for active psoriatic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Number of swollen joints
2. Number of tender joints
3. Dactylitis
4. Enthesitis
5. Skin and/or nail involvement

C. Active ankylosing spondylitis (AS) and active axial spondyloarthritis

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for active ankylosing spondylitis or active axial spondyloarthritis and who achieve or maintain a positive clinical response with the requested medication as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Functional status
2. Total spinal pain
3. Inflammation (e.g., morning stiffness)

D. Moderately to severely active ulcerative colitis (UC)

1. Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active ulcerative colitis and who achieve or maintain remission.
2. Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active ulcerative colitis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
 - i. Stool frequency
 - ii. Rectal bleeding
 - iii. Urgency of defecation
 - iv. C-reactive protein (CRP)
 - v. Fecal calprotectin (FC)
 - vi. Endoscopic appearance of the mucosa
 - vii. Improvement on a disease activity scoring tool (e.g., Ulcerative Colitis Endoscopic Index of Severity [UCEIS], Mayo score)

V. OTHER

For all indications: Member has had a documented negative TB test (which can include a tuberculosis skin test [PPD], an interferon-release assay [IGRA], or a chest x-ray)* within 6 months of initiating therapy for persons who are naïve to biologic DMARDs or targeted synthetic DMARDs associated with an increased risk of TB, and repeated yearly for members with risk factors** for TB that are continuing therapy with biologics.

* If the screening testing for TB is positive, there must be further testing to confirm there is no active disease. Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

** Risk factors for TB include: Persons with close contact to people with infectious TB disease; persons who have recently immigrated from areas of the world with high rates of TB (e.g., Africa, Asia, Eastern Europe, Latin America, Russia); children less than 5 years of age who have a positive TB test; groups with high rates of TB transmission (e.g., homeless persons, injection drug users, persons with HIV infection); persons who work or reside with people who are at an increased risk for active TB (e.g., hospitals, long-term care facilities, correctional facilities, homeless shelters).

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug.

VI. DOSAGE AND ADMINISTRATION

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

VII. APPENDICES

Appendix A: Examples of Contraindications to Methotrexate

1. Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease
2. Breastfeeding
3. Blood dyscrasias (e.g., thrombocytopenia, leukopenia, significant anemia)
4. Elevated liver transaminases
5. History of intolerance or adverse event
6. Hypersensitivity
7. Interstitial pneumonitis or clinically significant pulmonary fibrosis
8. Myelodysplasia
9. Pregnancy or currently planning pregnancy
10. Renal impairment
11. Significant drug interaction

Appendix B: Examples of Conventional Therapy Options for UC

1. Mild to moderate disease – induction of remission:
 - a. Oral mesalamine (e.g., Asacol, Asacol HD, Lialda, Pentasa), balsalazide, olsalazine
 - b. Rectal mesalamine (e.g., Canasa, Rowasa)
 - c. Rectal hydrocortisone (e.g., Colocort, Cortifoam)
 - d. Alternatives: prednisone, azathioprine, mercaptopurine, sulfasalazine
2. Mild to moderate disease – maintenance of remission:
 - a. Oral mesalamine, balsalazide, olsalazine, rectal mesalamine
 - b. Alternatives: azathioprine, mercaptopurine, sulfasalazine
3. Severe disease – induction of remission:
 - a. Prednisone, hydrocortisone IV, methylprednisolone IV

- b. Alternatives: cyclosporine IV, tacrolimus, sulfasalazine
- 4. Severe disease – maintenance of remission:
 - a. Azathioprine, mercaptopurine
 - b. Alternative: sulfasalazine
- 5. Pouchitis: Metronidazole, ciprofloxacin
 - a. Alternative: rectal mesalamine

VIII. REFERENCES

1. Simponi [package insert]. Horsham, PA: Janssen Biotech, Inc.; September 2019.
2. Van der Heijde D, Ramiro S, Landewe R, et al. 2016 Update of the international ASAS-EULAR management recommendations for axial spondyloarthritis. *Ann Rheum Dis.* 2017;76(6):978-991.
3. Smolen JS, Landewé RBM, Bijlsma JWJ, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis.* 2020;79(6):685-699. doi:10.1136/annrheumdis-2019-216655.
4. Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Rheumatol.* 2016;68(1):1-26.
5. Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis Rheum.* 2008;59(6):762-784.
6. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol.* 2011;65(1):137-174.
7. Gossec L, Baraliakos X, Kerschbaumer A, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies; 2019 update. *Ann Rheum Dis.* 2020;79(6):700-712.
8. Gladman DD, Antoni C, P Mease, et al. Psoriatic arthritis: epidemiology, clinical features, course, and outcome. *Ann Rheum Dis* 2005;64(Suppl II):ii14–ii17.
9. Peluso R, Lervolino S, Vitiello M, et al. Extra-articular manifestations in psoriatic arthritis patients. 2015;34(4):745-53.
10. Braun J, van den Berg R, Baraliakos X, et al. 2010 update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis. *Ann Rheum Dis.* 2011;70:896–904.
11. Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol.* 2019;71(10):1599-1613. doi:10.1002/art.41042.
12. Talley NJ, Abreu MT, Achkar J, et al. An evidence-based systematic review on medical therapies for inflammatory bowel disease. *Am J Gastroenterol.* 2011;106(Suppl 1):S2-S25.
13. Tuberculosis (TB). TB risk factors. Centers for Disease Control and Prevention. Retrieved on June 3, 2021 from: <https://www.cdc.gov/tb/topic/basics/risk.htm>.
14. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheumatol.* 2019;71(1):5-32. doi:10.1002/art.40726.
15. Feuerstein JD, Isaacs KL, Schneider Y, et al. AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. *Gastroenterology* 2020;158:1450.
16. Aletaha D, Neogi T, Silman, et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum.* 2010;62(9):2569-81.
17. Rubin DT, Ananthakrishnan AN, et al. 2019 ACG Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroentrol.* 2019;114:384-413.

Reference number(s)
2014-A

18. Smolen JS, Aletaha D. Assessment of rheumatoid arthritis activity in clinical trials and clinical practice. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. Available with subscription. URL: www.uptodate.com. Accessed March 19, 2021.
19. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthrit Care Res.* 2021;0:1-16.